



Rat Pulmonary Alveolar Epithelial Cells (RPAEpiC) Catalog #R3200

Cell Specification

Pulmonary alveolar epithelial cells, comprised of alveolar type I and type II epithelial cells, line more than 99% of the internal surface area of the lung [1]. Type I cells are large squamous cells whose thin cytoplasmic extensions cover >95% of the internal lung surface. They contain aquaporins and exhibit the highest osmotic water permeability of any mammalian cell type. Type II cells, which cover 2-5% of the internal lung surface, produce, secrete, and recycle pulmonary surfactant [2]. The currently accepted hypothesis is that Type II cells maintain pulmonary fluid homeostasis by regulating active Na⁺ transport in the lungs, whereas Type I cells are "inert" cells that provide only a barrier function. A recent study, however, suggests that Type I cells may also regulate ion and fluid transport [3].

RPAEpiC from ScienCell Research Laboratories are isolated from postnatal day 2 rat lung. RPAEpiC are cryopreserved at P0 and delivered frozen. Each vial contains >1 x 10⁶ cells in 1 ml volume. RPAEpiC are characterized by immunofluorescence with antibodies specific to CK-18, -19, and vimentin. RPAEpiC are negative for, mycoplasma, bacteria, yeast, and fungi. RPAEpiC are guaranteed to further culture under the conditions provided by ScienCell Research Laboratories. *However, this cell type is not recommended for expanding or long term cultures since the cells would differentiate to type I alveolar epithelial cells immediately after plating and type I alveolar epithelial cells do not proliferate in culture.*

Recommended Medium

It is recommended to use Alveolar Epithelial Cell Medium (AEpiCM, Cat. #3201) for culturing RPAEpiC *in vitro*.

Product Use

RPAEpiC are for research use only. They are not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

Storage

Upon receiving, directly and immediately transfer the cells from dry ice to liquid nitrogen and keep the cells in liquid nitrogen until they are needed for experiments.

Shipping

Dry ice.

References

- [1] Crapo JD, Young SL, Fram EK, Pinkerton KE, Barry BE, Crapo RO. (1983) "Morphometric characteristics of cells in the alveolar region of mammalian lungs." *Am Rev Respir Dis.* 128: S42-6.
- [2] Wright JR, Dobbs LG. (1991) "Regulation of pulmonary surfactant secretion and clearance." *Annu Rev Physiol.* 53: 395-414.
- [3] Johnson MD, Widdicombe JH, Allen L, Barbry P, Dobbs LG. (2002) "Alveolar epithelial type I cells contain transport proteins and transport sodium, supporting an active role for type I cells in regulation of lung liquid homeostasis." *Proc Natl Acad Sci USA.* 99: 1966-71.

Instructions for culturing cells

Caution: Cryopreserved cells are very delicate. Thaw the vial in a 37°C water bath and return the cells to culture as quickly as possible with minimal handling!

Initiating the culture:

1. Prepare a poly-L-lysine-coated culture vessel (2 $\mu\text{g}/\text{cm}^2$, T-75 flask is recommended). Add 10 ml of sterile water to a T-75 flask and then add 15 μl of poly-L-lysine stock solution (10 mg/ml, Cat. #0413). Leave the vessel in a 37°C incubator overnight (or for a minimum of one hour).
2. Prepare complete medium. Decontaminate the external surfaces of medium bottle and medium supplement tubes with 70% ethanol and transfer them to a sterile field. Aseptically transfer supplement to the basal medium with a pipette. Rinse the supplement tube with medium to recover the entire volume.
3. Rinse the poly-L-lysine-coated vessel twice with sterile water and then add 15 ml of complete medium. Leave the vessel in the sterile field and proceed to thaw the cryopreserved cells.
4. Place the frozen vial in a 37°C water bath. Hold and rotate the vial gently until the contents completely thaw. Promptly remove the vial from the water bath, wipe it down with 70% ethanol, and transfer it to the sterile field.
5. Carefully remove the cap without touching the interior threads. Gently resuspend and dispense the contents of the vial into the equilibrated, poly-L-lysine-coated culture vessel. A seeding density of 5,000 cells/ cm^2 is recommended.

Note: Dilution and centrifugation of cells after thawing are not recommended since these actions are more harmful to the cells than the effect of residual DMSO in the culture. It is also important that cells are plated in poly-L-lysine-coated culture vessels to promote cell attachment.

6. Replace the cap or lid of the culture vessel and gently rock the vessel to distribute the cells evenly. Loosen cap, if necessary, to allow gas exchange.
7. Return the culture vessel to the incubator.
8. For best results, do not disturb the culture for at least 16 hours after the culture has been initiated. Refresh culture medium the next day to remove residual DMSO and unattached cells, then every other day thereafter.

Maintaining the culture:

1. Refresh supplemented culture medium the next morning after establishing a culture from cryopreserved cells.
2. Change the medium every two to three days thereafter.

It is not recommended that RPAEpiC be subcultured beyond their initial plating.

Caution: Handling animal-derived products is potentially biohazardous. Always wear gloves and safety glasses when working with these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination [1].

[1] Grizzle WE, Polt S. (1988) "Guidelines to avoid personal contamination by infective agents in research laboratories that use human tissues." *J Tissue Cult Methods*. 11: 191-9.